Arcadia Planitia: Acheron Fossae and Erebus Montes Workshop Abstract #1011


Donna Viola¹ (dviola@lpl.arizona.edu), Alfred S. McEwen¹, & Colin M. Dundas²

¹University of Arizona, Department of Planetary Sciences, ²USGS Astrogeology Science Center

Motivation: Shallow Subsurface Water Ice



 $1^{
m st}$ EZ Workshop for Human Missions to Mars



70° N

60° N

50° N

110° W - 70°

60° N

Cluster Type

Mixed

Expanded Normal

Expanded Secondary Craters

These craters require substantial excess ice in order to form. They suggest that the Arcadia Planitia ice sheet is tens of millions of years old.

Viola, et al. (2015). Icarus 248: 190-204.


Arcadia Planitia is one of the few regions where abundant shallow ice is present at relatively low latitude.

Motivation: Shallow Subsurface Water Ice



 $1^{
m st}$ EZ Workshop for Human Missions to Mar $\,$

Map of recent ice-exposing impact craters




300

240

30°

Dust cover index basemap (red = dusty, blue = less dusty)



-60

180

120°

These craters expose "excess" ice (Dundas et al., 2015)— which is almost entirely free of dust (>99% water ice).

Water-equivalent hydrogen basemap (blue = more ice, orange = little/no ice)


Arcadia Planitia is one of the few regions where abundant shallow ice is present at relatively low latitude.

Motivation: Shallow Subsurface Water Ice



 $1^{
m st}$ EZ Workshop for Human Missions to Mar $\,$

Map of recent ice-exposing impact craters




300

240

-30'

Dust cover index basemap (red = dusty, blue = less dusty)



-60

180

120°


These craters expose "excess" ice (Dundas et al., 2015)— which is almost entirely free of dust (>99% water ice).

Water-equivalent hydrogen basemap (blue = more ice, orange = little/no ice)

Arcadia Planitia is one of the few regions where abundant shallow ice is present at relatively low latitude.


Arcadia Planitia: Geologic Context





Arcadia Planitia: Geologic Context







Arcadia Planitia: Challenges



1st EZ Workshop for Human Missions to Mars





Acheron Fossae


Erebus Montes


Dust: Arcadia Planitia is fairly dusty. Landing sites were chosen in areas that appear to minimize dust using the TES Dust Cover Index.

Arcadia Planitia: Challenges



1st EZ Workshop for Human Missions to Mars





Acheron Fossae


Erebus Montes


Composition: Very little CRISM data for either EZ.

Arcadia Planitia: Challenges



1st EZ Workshop for Human Missions to Mars






Acheron Fossae

Erebus Montes

High-resolution coverage: Some HiRISE coverage, but not much.




Donna Viola¹ (dviola@lpl.arizona.edu), Alfred S. McEwen¹, & Colin M. Dundas²

¹University of Arizona, Department of Planetary Sciences, ²USGS Astrogeology Science Center


Exploration Zone Map



1st EZ Workshop for Human Missions to Mars




SCIENCE ROIS






1st EZ Workshop for Human Missions to Mars





- 38.7N, 221E. Well-covered in CTX, minimal HiRISE.
- Acheron Fossae graben system, late Noachian highlands terrain.
 - Presence of Noachian rocks.
 - Unit with regional/global extent.
 - Significant tectonic activity.
- Many different features of interest within this larger region.








12














- Exposures along graben walls (multiple locations).
 - Access to Noachian bedrock units?












- Glacial fill on graben floors (likely Amazonian in age).
 - Range of surface ages (relative).
 - Access to Amazonian subsurface ice.







- 38.9N, 219.5E. Covered by CTX, HiRISE.
- Small channel-like feature, not in fluvial network – possible evidence of sub-glacial liquid water flow.
 - Aqueous activity
 - Potential habitability?






7.5 Kilometers


- 40.8N, 219.6E. CTX coverage, little HiRISE.
- Periglacial-type terrain; large blocky mesas
- Abundant subsurface ice (Amazonian), including glacial fill between blocks.













- 40.3N, 221E. CTX coverage, some HiRISE.
- Double layer ejecta impact crater, concentric crater fill, radial texture apparent in parts of ejecta.
 - Access to
 Amazonian aged
 subsurface ice.
 - Primary crater deposits (with unique ejecta morphology).

RESOURCE ROIs



Resource ROIs





- RROI-1: 39.8N, 220.8E
 RROI-2: 39.7N, 220.6E
- Sample RROI locations near HZ, on the fringes of lobate, likely icerich, ejecta material.
- Excess water ice thought to be abundant throughout this entire region.

Science ROI(s) Rubric



1st EZ Workshop for Human Missions to Mars

			Site Factors	SR011	SR012	SROI3	SR014	RR011	RR012	EZ SUM	
	oio	Threshold	Potential for past habitability	0	0	?	?	?	?	0,2	
	Astrobio	Till Calloid	Potential for present habitability/refugia	0	0	0	0	0	0	0,6	
	Ä	Qualifying	Potential for organic matter, w/ surface exposure	?	?	?	?	?	?		
	nce	Threshold	Noachian/Hesperian rocks w/ trapped atmospheric gases	0	0					0,2	
	Scie		Meteorological diversity in space and time	?	?	?	?	?	?		
	Atmospheric Science	0 115	High likelihood of surface-atmosphere exchange	•	•	•	•	•	•	6,0	
Site Criteria	losph	Qualifying	Amazonian subsurface or high-latitude ice or sediment	•	•	•	•	•	•	6,0	
Crit	Atm		High likelihood of active trace gas sources	?	?	?	?	?	?		
te (Range of martian geologic time; datable surfaces	•		•				2,0	
		Threshold	Evidence of aqueous processes	•	•					2,0	
nce			Potential for interpreting relative ages	•	?	?	•			2,0	
Science	ce		Igneous Rocks tied to 1+ provinces or different times	?	?	?	?	?	?		
•	cien		Near-surface ice, glacial or permafrost	•	•	•	•	•	•	6,0	
	Geoscience		Noachian or pre-Noachian bedrock units	•	•					2,0	
)	Qualifying	Outcrops with remnant magnetization	?	?	?	?	?	?		
			Primary, secondary, and basin-forming impact deposits	•	•	•	•			4,0	
			Structural features with regional or global context	•						1,0	
			Diversity of aeolian sediments and/or landforms	?	?	?	?	?	?		

Key							
•	Yes						
0	Partial Support or Debated						
	No						
?	Indeterminate						

Resource ROI(s) Rubric




Lst EZ Workshop for Human Missions to Mar

		Site Factors	SROI	SROL	SROI	SROL	RROI	RROL	EZ SU
E	ngineering	Meets First Order Criteria (Latitude, Elevation, Thermal Inertia)	•	•	•	•	•	•	6,0
		Potential for ice or ice/regolith mix	•	•	•	•	•	•	6,0
		Potential for ice or ice/regolith mix Potential for hydrated minerals	?	?	?	?	?	?	
l e		Quantity for substantial production	•	•	•	•	•	•	6,0
l j	Threshold	Potential to be minable by highly automated systems	•	•	•	•	•	•	6,0
Res		Located less than 3 km from processing equipment site					•	•	2,0
Water Resource		Located no more than 3 meters below the surface	•	•	•	•	•	•	6,0
Wat		Accessible by automated systems					•	•	2,0
		Potential for multiple sources of ice, ice/regolith mix and hydrated minerals							
)	Qualifying	Distance to resource location can be >5 km							
_		Route to resource location must be (plausibly) traversable							
ng		~50 sq km region of flat and stable terrain with sparse rock distribution							
eeri	Threshold	1–10 km length scale: <10°	0	0	•	0	•	•	3,3
Civil Engineering		Located within 5 km of landing site location					•	•	2,0
ᆸ		Located in the northern hemisphere	•	•	•	•	•	•	6,0
	Qualifying	Evidence of abundant cobble sized or smaller rocks and bulk, loose regolith	?	?	?	?	?	?	
ت ا		Utilitarian terrain features	?	?	?	?	?	?	
5	5	Low latitude							
اج ال	Š	No local terrain feature(s) that could shadow light collection facilities							
Food	Qualifying	Access to water	•	•	•	•	•	•	6,0
ءُ ا		Access to dark, minimally altered basaltic sands	?	?	?	?	?	?	
		Potential for metal/silicon	?	?	?	?	•	•	2,0
٦		Potential to be minable by highly automated systems					?	?	
	Threshold	Located less than 3 km from processing equipment site					•	•	2,0
Metal/Silicon		Located no more than 3 meters below the surface					0	0	0,2
eta		Accessible by automated systems					?	?	
Σ̈́		Potential for multiple sources of metals/silicon							
	Qualifying	Distance to resource location can be >5 km							
		Route to resource location must be (plausibly) traversable							

ISRU and Civil Engineering Criteria

Key								
•	Yes							
0	Partial Support or Debated							
	No							
?	Indeterminate							




Donna Viola¹ (dviola@lpl.arizona.edu), Alfred S. McEwen¹, & Colin M. Dundas²

¹University of Arizona, Department of Planetary Sciences, ²USGS Astrogeology Science Center

Exploration Zone Map




1st EZ Workshop for Human Missions to Mars




Erebus Montes 23

SCIENCE ROIS


Erebus Montes 24








- 39.7N, 190.6E. Well-covered in CTX, some HiRISE.
- Exposures of Hesperian-Noachian transition terrain.
 - Noachian/Hesperian rocks?
- Some mounds of HNt material have ice-rich lobate debris aprons.
 - Amazonian-aged subsurface ice.
 - Relative ages.
- Located near a recent ice-exposing impact crater.
 - Subsurface ice.
 - Impact crater processes.




Erebus Montes


25



1st EZ Workshop for Human Missions to Mars




- 39.5N, 194E. Well-covered in CTX, some HiRISE.
- Additional exposures of Hesperian-Noachian transition terrain.
- Some mounds of HNt material have ice-rich lobate debris aprons. Filled/buried crater seen in SE region.
 - Impact processes, relative ages.










- 38.5N, 192.8E. Wellcovered in CTX, no HiRISE.
- Contact between multiple episodes of Amazonian volcanism.
 - Relative surface ages.
 - Possibility for lava tube caves?
- Also includes some evidence for subsurface ice.

27 Frebus Montes









- 39.5N, 191.5E. Well-covered in CTX, no HiRISE.
- Filled/buried impact crater, evidence for glacial/ periglacial processes.



Erebus Montes 28



1st EZ Workshop for Human Missions to Mars



- 37.9N, 190.8E. Well-covered in CTX, some HiRISE.
- Mounds with lobate debris aprons, surrounded by Amazonian lava flows. Evidence for glacial/


RESOURCE ROIS

Erebus Montes 30



Resource ROIs





RROI-1: 39.1N, 192E
 RROI-2: 38.9N, 192.2E

 Sample RROI locations near HZ. Both contain evidence for excess subsurface water ice (thought to be abundant throughout this entire region).

Erebus Montes

Science ROI(s) Rubric

1st EZ Workshop for Human Missions to Mars

				Site Factors	SR011	SROI2	SROI3	SR014	SROIS	RR011	RR012	EZ SUM
	oio	Threshold	J/OR	Potential for past habitability	?	?	?	?	?	?	?	
	Astrobio	Tillesiloid	AND	Potential for present habitability/refugia	0	0	0	0	0	0	0	0,7
	As	Qualifying		Potential for organic matter, w/ surface exposure	?	?	?	?	?	?	?	
	nce	Threshold	Noa	chian/Hesperian rocks w/ trapped atmospheric gases	0	0			0			0,3
	Scie			Meteorological diversity in space and time	?	?	?	?	?	?	?	
	Atmospheric Science	0 115		High likelihood of surface-atmosphere exchange	•	•	•	•	•	•	•	7,0
Criteria	nosph	Qualifying		Amazonian subsurface or high-latitude ice or sediment	•	•	•	•	•	•	•	7,0
Crit	Atm			High likelihood of active trace gas sources								
te				Range of martian geologic time; datable surfaces	•	•	•	•	•			5,0
_		Threshold		Evidence of aqueous processes								
D (I)		Inresnoia										1
nce s		Inresnoia		Potential for interpreting relative ages	•	•	•	•	•			5,0
science s	се	inresnoia		Potential for interpreting relative ages Igneous Rocks tied to 1+ provinces or different times	?	•	•	• ?	• ?	?	?	5,0
Science S	cience	Inresnoid			•		?	?		?	?	5,0 7,0
Science Site	Seoscience	Inresnoid		Igneous Rocks tied to 1+ provinces or different times	• ? • 0	?		?	?		?	-
Science S	Geoscience	Qualifying		Igneous Rocks tied to 1+ provinces or different times Near-surface ice, glacial or permafrost	•	?		?	?		?	7,0
Science S	Geoscience			Igneous Rocks tied to 1+ provinces or different times Near-surface ice, glacial or permafrost Noachian or pre-Noachian bedrock units	•	?	•	•	?	•	•	7,0
Science S	Geoscience			Igneous Rocks tied to 1+ provinces or different times Near-surface ice, glacial or permafrost Noachian or pre-Noachian bedrock units Outcrops with remnant magnetization	•	?	•	•	?	•	•	7,0 0,3

Key							
•	Yes						
0	Partial Support or Debated						
	No						
?	Indeterminate						

Resource ROI(s) Rubric



1st EZ Workshop for Human Missions to Mars

			Site Factors	SRO]	SRO	SRO]	SRO]	SRO	RRO	RRO	EZ SI
	En	gineering	Meets First Order Criteria (Latitude, Elevation, Thermal Inertia)	•	•	•	•	•	•	•	7,0
			Potential for ice or ice/regolith mix	•	•	•	•	•	•	•	7,0
			Potential for hydrated minerals	?	?	?	?	?	?:	?	
	rce		Quantity for substantial production	•	•	•	•	•	•	•	7,0
	our	Threshold	Potential to be minable by highly automated systems	•	•	•	•	•	•	•	7,0
ia	Resource		Located less than 3 km from processing equipment site						•	•	2,0
Criteria	er F		Located no more than 3 meters below the surface	•	•	•	•	•	•	•	7,0
rit	Water		Accessible by automated systems						•	•	2,0
J	_		Potential for multiple sources of ice, ice/regolith mix and hydrated minerals								\Box
9		Qualifying	Distance to resource location can be >5 km								
in			Route to resource location must be (plausibly) traversable								
er	ηg		~50 sq km region of flat and stable terrain with sparse rock distribution								
and Civil Engineering	Engineering	Threshold	1-10 km length scale: <10°	0	•	•	0	•	•	•	5,2
gir	jine		Located within 5 km of landing site location						•	•	2,0
u:	Eng		Located in the northern hemisphere	•	•	•	•	•	•	•	7,0
3	Civil	Qualifying	Evidence of abundant cobble sized or smaller rocks and bulk, loose regolith	?	?	?	?	?	?	?	
Vi	ί		Utilitarian terrain features		?	?	?	?	?	?	
Ö	n		Low latitude								
<u> </u>	الظوا		No local terrain feature(s) that could shadow light collection facilities								
Ĕ	Food Production	Qualifying	Access to water	•	•	•	•	•	•	•	7,0
5 L			Access to dark, minimally altered basaltic sands	?	?	?	?	?	?	?	
ISRU	rce		Potential for metal/silicon	?	?	?	?	?	•	•	2,0
SI	Resource		Potential to be minable by highly automated systems						•	•	2,0
		Threshold	Located less than 3 km from processing equipment site						•	•	2,0
	Metal/Silicon		Located no more than 3 meters below the surface						0	0	0,2
	ili		Accessible by automated systems						?	?	
	5/1		Potential for multiple sources of metals/silicon								
	eta	Qualifying	Distance to resource location can be >5 km								
	Σ̈́		Route to resource location must be (plausibly) traversable								

Key								
•	Yes							
0	Partial Support or Debated							
	No							
?	Indeterminate							

BONUS Science ROI



1st EZ Workshop for Human Missions to Ma

For both Acheron Fossae and Erebus Montes: The Arcadia Planitia Ice Sheet

- Easy access to subsurface ice means that it should be easy to sample and study as well.
- Science questions:
 - How did the ice sheet form?
 - How long did the ice deposition take, and how long has it existed?
 - How thick is the ice sheet?
 - How is the ice interacting with the atmosphere?
 - Did the ice preserve organics or other biosignatures?
 - Could the ice have contained habitable microenvironments now or in the past?



1st EZ Workshop for Human Missions to Ma

For both Acheron Fossae and Erebus Montes: The Arcadia Planitia Ice Sheet

- Easy access to subsurface ice means that it should be easy to sample and study as well.
- Science questions:
 - How did the ice sheet form?
 - How long did the ice deposition take, and how long has it existed?
 - How thick is the ice sheet?
 - How is the ice interacting with the atmosphere?
 - Did the ice preserve organics or other biosignatures?
 - Could the ice have contained habitable microenvironments now or in the past?

Water, water (ice) everywhere – but not all of it's to drink!



1st EZ Workshop for Human Missions to Ma

- Science Priority: surface composition data
 - Surface dust affects CRISM observations.
 - Strategy: Target CRISM observations in regions with slightly less dust.
- Resource Priority: higher resolution of nearsurface ice deposits, including spatial distribution
 - SAR RADAR, higher resolution neutron spectrometer